

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of selection for scrapie resistance on genetic diversity in a rare and locally adapted sheep breed: The case of Sambucana.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/139173> since

Published version:

DOI:10.1016/j.livsci.2013.08.006

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in

Effect of selection for scrapie resistance on genetic diversity in a rare and locally adapted sheep breed: The case of Sambucana.

Sartore, S., Rasero, R., Colussi, S., Acutis, P.L., Peletto, S., Soglia, D., Maione, S., Spalenza V., Sacchi, P. (2013) *Livestock Science* 157 (1) PP. 75 - 80 doi: 10.1016/j.livsci.2013.08.006

<http://www.livestockscience.com/article/S1871-1413%2813%2900355-7/abstract>

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>),
<http://www.sciencedirect.com/science/article/pii/S1871141313003557>

Effect of selection for scrapie resistance on genetic diversity in a rare and locally adapted
sheep breed: the case of Sambucana

S. Sartore¹, R. Rasero¹, S. Colussi², P.L. Acutis², S. Peletto², D. Soglia¹, S. Maione¹, V.
Spalenza¹, P. Sacchi¹

¹Dipartimento di Produzioni Animali Epidemiologia ed Ecologia, via Leonardo da Vinci 44
Grugliasco (Torino), Italy

²Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle d'Aosta, via Bologna
148, 10154 Torino, Italy

Correspondence

Paola Sacchi, Dipartimento di Produzioni Animali Epidemiologia ed Ecologia, via
Leonardo da Vinci 44 Grugliasco (Torino), Italy

Tel.: 0039 (0)11 6709254,

Fax: 0039 (0)11 6709240

E-mail paola.sacchi@unito.it

Abstract.

In Italy, a breeding plan has been adopted in 2005 to increase resistance to scrapie. The effect of selection on genetic diversity of Sambucana, a rare and locally adapted sheep breed, was assessed by analysing the evolution of allele frequencies at different levels: the PRNP (prion protein) gene itself, microsatellites on the same chromosome as PRNP, and microsatellites on other chromosomes, not subjected to selection for resistance to scrapie. A total of 147 young rams, 80 born in 2004 and 67 born in 2008–2009 were analysed. Evidence of diversity loss was observed for PRNP as a consequence of the directional selection. Diversity was affected in the immediate vicinity of PRNP but the effect on more distant loci on the same chromosome was trivial. With regard to neutral markers, lack of heterozygosity with no changeover of allele frequencies was observed suggesting an increase of inbreeding. Mating policies would be sufficient to solve these problems. A selection scheme based on genotyping rams and eliminating carriers of susceptibility and all carriers of high susceptibility is the best way to improve natural resistance to scrapie with low costs and minimal problems in the current conservation programmes targeting rare breeds.

Keywords: sheep; scrapie resistance; genetic variation; prion protein gene; molecular marker.

46 Introduction.

47 In sheep, susceptibility to scrapie is influenced by mutations in the coding region of the
48 PRNP (prion protein) gene, located on OAR13 (sheep chromosome 13). Haplotypes-alleles
49 valine/arginine/glutamine (VRQ) and alanine/arginine/glutamine (ARQ) at codons 136,
50 154, and 171, respectively, are associated with high susceptibility, whereas the
51 alanine/arginine/arginine allele (ARR) has been linked to decreased susceptibility or
52 resistance (Belt et al., 1995; Bossers et al., 1996; Hunter et al., 1996).

53 Accordingly, the European Union has implemented programmes for genetic control
54 of scrapie susceptibility in sheep (European Commission, 2003). Since 2005 in Italy, a
55 breeding plan has been adopted to increase the frequency of the ARR ‘resistant’ allele and
56 to eliminate the VRQ ‘susceptible’ allele (Decreto Ministeriale 17 Dicembre 2004).

57 The impact of selection for scrapie resistance has been investigated frequently.
58 Different selection strategies applied to breeds with different genetic structure were
59 considered. Most studies predict effects and costs of selection in term of variability using
60 both simulated and real datasets (Alfonso et al., 2006; Álvarez et al., 2007; Álvarez et al.,
61 2009; Drögemüller et al., 2004; Man et al., 2009; Molina et al., 2006; Roden et al., 2006;
62 Windig et al., 2004; Wiśniewska et al., 2010), whereas only a few investigation analyse the
63 realized effects after the selection programmes have been applied (Palhière et al., 2006;
64 Palhière et al., 2008). Various concerns have been raised regarding possible unintended
65 consequences of widespread selection on PRNP alleles, including risk of genetic diversity
66 loss and increased inbreeding (Dawson et al., 2008; Parada et al., 2007).

67 The Sambucana is a Piemonte region breed devoted to meat production. Since 1985,
68 several initiatives have been carried out to safeguard the breed and economically valorise
69 derived productions. An important step in the conservation programme was the creation of

the 'Agnello Sambucano Garantito' brand, which certifies the origin of the lambs and guarantees the quality of the meat. In 2001, the Sambucano lamb was added to the 'Presidia' list of Slow Food. In 1993 the breed was classified by the FAO as 'at limited diffusion' (FAO-UNEP, 1993). In 2005, 168 rams and 3995 ewes were registered (Associazione Nazionale della Pastorizia, <http://www.assonapa.it>).

The Sambucana breed showed an ARR frequency above the threshold to comply with European regulations, before the selection programme started. Nevertheless, as a rare and locally adapted breed, it could benefit of some derogations. In Piemonte, the selection programme for scrapie resistance has been applied since 2005 (Regione Piemonte, 2005). This decision meets consumer expectations of a high-quality meat that is also safe to eat.

Currently, great attention is paid to avoid excessive inbreeding or genetic drift. Genealogical information can be used to monitor the evolution of genetic variability. However, in a population kept in an extensive breeding system, the quality of pedigree information available is often inadequate. For the Sanbucana, pedigrees are not available at all. In such cases, the evolution of genetic variability can be assessed using a molecular approach.

The aim of the present investigation was to use Sambucana as a model to evaluate the consequences of selection for scrapie resistance on molecular variability in a rare and locally adapted breed. Three processes may play a role affecting genetic variability: direct selection on PRNP itself, frequency changes in marker loci on the same chromosome because of the linkage, and genetic drift because a limited part of the population is used as parents. Therefore, the evolution of allele frequencies was analysed at these different levels: PRNP gene, microsatellite marker loci on OAR13, and microsatellites on chromosomes other than OAR13, markers not subjected to direct selection for scrapie resistance.

94

95 Materials and methods.

96 Sample collection. A total of 147 Sambucana young rams (candidate sires), 80 born in 2004
97 and 67 born in 2008–2009 (denoted as the before-2005 and after-2005 cohorts), were
98 randomly chosen from different flocks among all animals genotyped at the PRNP locus by
99 the IZSTO-CEA (Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle
100 d'Aosta, Italian Reference Centre for Animal Transmissible Spongiform
101 Encephalopathies). The between-sampled cohorts period length (4 years) represents about
102 one generation.

103 The analyses were performed on rams only because, before 2005, males were
104 genotyped exclusively.

105 Each cohort was divided into two risk groups according to the selection criteria
106 adopted in the Italian breeding plan against VRQ and in favour of ARR. 'Low risk' rams
107 are ARR/ARR and ARR/non-ARR, except ARR/VRQ, all other animals being considered
108 as 'high risk'.

109 Molecular techniques. Genomic DNA was extracted from blood samples using the
110 NucleoSpin QuickPure extraction kit (Macherey-Nagel, Dueren, Germany). Fourteen
111 microsatellites were chosen on OAR13 (Table 1). Five markers are strictly linked to PRNP:
112 PRNPS11, -15, and -24 map within PRNP whereas PRNPS04 and -05 map at about 40 kb
113 upstream of the 5' end of the gene sequence GenBank U67922.1. The other nine OAR13
114 markers were chosen outside the gene at various distances to verify whether the effect of
115 selection depends on relative distances (Geldermann et al., 2003; Isler et al., 2006; Lühken
116 et al., 2006; Palhière et al., 2008; Preuss et al., 2005). Other 12 microsatellites, belonging to
117 the ISAG (<http://www.isag.us/>) and the ECONOGENE (<http://www.econogene.eu/>)

recommended panels, were selected to avoid syteny with PRNP (denoted as neutral
microsatellites in Table 1). PCR assays were developed and performed as previously
described (Soglia et al., 2010).

In order to validate the microsatellite panel, an error assay was performed by
replicating the genotyping molecular analyses on a randomly chosen 10% of individual
samples. The average error rate per allele was computed as suggested by Pompanon et al.
(2005). Individual markers in each age cohort were tested for deviations from HW (Hardy–
Weinberg) proportions, F_{IS} statistics and the significance of their non-zero values were
obtained using Fstat 2.9.3.2 software (Goudet, 1995). The presence of linkage
disequilibrium between neutral markers was checked using the likelihood ratio test
implemented with Arlequin software (Excoffier et al., 2005).

Analyses of genetic variation. Genetic variation was analysed for PRNP and for the
markers grouped into three clusters: OAR13 microsatellites within PRNP, OAR13
microsatellites outside PRNP, and neutral microsatellites. Departures from the HW
proportions were examined using the F_{IS} statistics.

Losses of both heterozygosis and number of alleles in term of within-groups,
between-groups, and total contributions to variability were assessed following Caballero
and Toro (2002) and Petit et al. (1998) approaches as proposed by Álvarez et al. (2009)
using MolKin 3.0 (Gutiérrez et al., 2005). Raw number of alleles (A), allelic richness, or
number of alleles adjusted for sampling size (A_g , g being two-fold the number of
individuals with full genotype information in the group of lowest size) (Hurlbert, 1971),
gene diversity (GD), or expected heterozygosis (Nei, 1987), and PIC (polymorphism
information content) were calculated. At both age cohorts, loss of genetic variability was
quantified sequentially removing all the ‘high risk’ rams.

At all 26 microsatellite loci, differences in allele frequencies between cohorts were tested with the F_{ST} statistics as implemented by the Arlequin software applying analysis of molecular variance.

Results.

Validation of the microsatellite markers. The rate of missing genotypes was 1.4%, and the average error rate per allele was 0.2%. Eleven individual samples were dropped from the dataset because they failed to consistently provide amplification products at all or most loci. The data for 26 microsatellites from 136 individual samples (71 and 65 from the two age cohorts, respectively) were finally analysed. Three markers revealed systematic deviations from HW proportions in both age cohorts whereas seven loci showed significant deviations only after 2005 (Table 1). The test for linkage disequilibrium revealed no significance among neutral microsatellites. These results characterized the chosen microsatellites as a useful tool to obtain the goals of our investigation.

Analyses of genetic variation of the PRNP gene. All five known PRNP alleles were observed (Table 2). ARQ was predominant before 2005 when ARH was the least frequent allele. After one generation of selection, ARR doubled in frequency, becoming the predominant allele, whereas ARQ dropped by 38% and VRQ by 69%. ARH was not sampled after 2005; consequently, number of alleles was reduced by one. The reversal of the frequency ratio between ARR and ARQ affected the genotypic frequencies and thus the availability of resistant rams. The percentage of ARR/ARQ rams was 43.7% in the before-2005 cohort and 38.5% one generation later; however, the ARR/ARR rams, representing only 2.8% before 2005, expanded to 35.4% in the later cohort. As a whole, the 'low-risk'

rams increased from 49% to 78%. The rates of ARQ/ARQ were 28.2% and 12.3% in the two age cohorts, respectively. The VRQ-carrier rams decreased from 12.7% to 4.6%.

Analyses of genetic variation at the microsatellite loci. The before-2005 and after-2005 cohorts were compared for their allelic arrangements (Table 3). The difference between the two age cohorts, as tested using the F_{ST} statistics, was highly significant for PRNP. Likewise, significant differences were identified at the OAR13 microsatellites for both within and outside PRNP markers. The neutral markers, on the other hand, showed no modification of their allele frequencies after one generation. In Figure 1 the differences for each OAR13 marker separately are plotted versus the distance to the PRNP locus.

Microsatellites within PRNP on the overall showed a decline in heterozygosis after the selection plan started (Table 4). The GD and PIC decreased between the two cohorts by 26% within PRNP, but by no more than 2% on the other locations. No decline in number of alleles (A and A_g) was observed on OAR13 whereas, after the selection plan started, the allelic richness increased by 7% at the neutral markers.

A detailed analysis of individual markers showed that PRNPS11, which is on PRNP, lost one allele and 59% of its original heterozygosis, whereas PRNPS04 lost 48% of its heterozygosis but no alleles. The other three markers within PRNP showed lower decreases in heterozygosis (16–17%). The OAR13 microsatellites outside PRNP showed only small differences between cohorts except BMS2319, that underwent a pronounced loss of diversity.

OAR13 markers outside PRNP after 2005 and neutral markers in both age cohorts showed observed heterozygosis deficiency, characterized by positive and significant F_{IS} values (Table 4).

These results show that the effects of selection on OAR13 microsatellites differed according to the relative distance of the loci from the PRNP gene and their polymorphism.

Changes in each of the two age cohorts were then computed after removal of all 'high-risk' rams to ascertain if overall contributions to variability were negligible or not (Table 5).

The cull of the 'high-risk' rams from both age cohorts dramatically decreased the GD_T within PRNP (more than 15%) with negative contributions mainly at GD_W . In fact, the 'high-risk' rams carried more heterozygosity than the selected ones (0.464 vs. 0.307 and 0.373 vs. 0.268 for the two age cohorts, respectively). Although to a small extent (less than 2%), the selection decreased the GD_T also at the OAR13 markers outside PRNP.

On the other genomic locations, selectively neutral, the lowest decrease of GD_T was observed before 2005, whereas in the succeeding generation the same parameter even showed a small increase.

The 'high-risk' rams provided a positive contribution to the allelic richness at the within PRNP markers, but only in the younger cohort. After removal of the same animals, on the OAR13 markers outside PRNP the allelic richness decreased at both within and total levels because the culled rams provided positive contributions. In spite of this, the overall allelic richness did not decrease after 2005 (Table 4).

On the contrary, the first selective action did not decrease diversity on the other chromosomes, where the 'high-risk' rams provided an unfavourable contribution; as a consequence, the cull of these animals gave rise to the gain in A and A_g observed in Table 4 between the two cohorts.

Discussion.

The main goal of the present analysis was to assess the effect of the selection for scrapie resistance on genetic variability in a small and locally adapted breed at the early stage of a selection programme. Genealogical information can be used to monitor the evolution of genetic diversity. Unfortunately, pedigrees were not available for this breed. Therefore, information from molecular investigation was the best option to assess the effect of selection on the genetic variability (Álvarez et al., 2009). Unlike simulation studies, based on genetic characteristics of a population, our investigation used real population data both before and after the selection acted. The Sambucana breed was used as a model.

Before the selection for scrapie resistance started, the predominant PRNP allele was ARQ. This allele is also predominant in most Italian and European breeds and is thought to represent the ancestral form of the PRNP gene (Acutis et al., 2003; Goldmann et al., 2005). From 2005 onwards, the selection scheme for scrapie resistance has provided for all ARR/ARR rams to be bred first and then ARR/non-ARR rams with equal preference, with ARR/VRQ rams being excluded.

Under the selection programme, despite the short time interval (one generation), evidence of variability loss was observed for PRNP as a consequence of the directional selection acting on this gene. In fact, in the most recent cohort, an allele had been lost. The change in genotypic frequencies illustrates the effectiveness of the selection programme. The ARR carriers reached a proportion of 78% in only a few years despite the fact that the ARR allele frequency was rather low at the beginning of selection. On the other hand, the ARQ and VRQ frequencies decreased significantly. Because the selection programme did not involve females, the VRQ allele could not be eliminated in a single generation. Overall, the implementation of a mild selection strategy achieved a satisfactory increase in ARR and a large decrease in VRQ. On the other hand, at the beginning of the selection plan

implementation, the number of homozygote rams was small, and the use of ARR/ARR males exclusively would have adversely affected the within-population genetic variability through a severe bottleneck. In the future, a changeover to using only homozygote rams will be possible to accelerate the increase in ARR frequency.

The selection changed also the variability of the five microsatellites located within the PRNP sequence. In particular, selection for ARR strongly affected PRNPS11. ARR was quite exclusively linked to the PRNPS11-151 allele, whereas the VRQ allele was exclusively linked to the 149 allele. Selection had less effect on PRNPS04, -05, -15, and -24, probably because of their low initial informative content. The effect of selection on OAR13 markers outside PRNP seemed to be quite low, with average differences between cohorts very close to zero. This positional relativity of effect has been identified previously; in four French breeds, Palhière et al. (2008) found that the effect on markers at OAR13 depended strongly on the relative distance from the selection objective. In Sambucana, selection affected genomic diversity in the immediate vicinity of PRNP, but the effect on more distant loci on the chromosome was trivial. The signature shown by BMS2319 was difficult to trace back to selection for resistance to scrapie because this microsatellite is the most distant from PRNP.

With regard to neutral markers, a lack of observed heterozygosis, with no changeover of allele frequencies, was noted in both age cohorts (Table 3 and 4). These findings suggest presence of inbreeding already before selection, due to the small size of the population, but the tendency is increasing. In this last respect, several explanations are possible: 1) because only 49% of young rams had a favourable genotype before 2005, the narrow number of selected reproducers reduced the effective population size still further; 2) in a small population, the carriers of favourable genotypes may be more related to each

other than randomly chosen individuals, and for an equal number of reproducers, the effective size thus may be smaller than expected in a pure genetic drift condition; 3) the farmers who owned young resistant rams could have used them intensively in their native flocks, reducing gene flow between flocks. Mating policies implemented to avoid inbreeding would be sufficient to solve these problems.

Except for the OAR13 markers, removal of susceptible rams did not reduce heterozygosis and allelic richness, either within a generation or across generations. This may be due to similar allele frequency distribution in selected as well as in culled rams. The genetic background of the 'high risk' rams was well represented in the selected group and, in addition, the cull of these animals gave rise to gain in number of alleles between the two generations. It must be noted that rams tested after 2005 were progeny of both selected rams and unselected ewes born before 2005. Therefore, they provided a sample of the allele pool of the parents that was achieved with the first selective decisions. Moreover, because the resistant rams born after 2005 also showed no reduction in allelic richness, it can be inferred that the population is not in danger of a strong bottleneck.

Conclusion.

Allelic richness and gene diversity are important for conservation of genetic stocks. The long-term evolutionary potential of a population and the limit of selection response are determined by the initial number of alleles as well as by the number of conserved alleles, regardless of the allele frequencies (Falconer and Mackay, 1996).

The consequences of adding a new criterion of selection differ between breeds depending on the initial situation and the strategy applied. The results based on the Sambucana breed can be generalized and the recommendation extended to other breeds

with similar starting genetic properties. The carriers of undesirable PRNP genotypes would not be essential to maintain genetic variability in the overall genome outside OAR13.

Selection scheme based on genotyping rams and eliminating non-ARR/non-ARR and all VRQ carriers is actually the best way to improve natural resistance to scrapie with low costs and minimal problems in the current conservation programme targeting rare breeds. The drawback of this strategy is that it does not result in the immediate elimination of VRQ; nevertheless, it still guarantees a rapid and considerable reduction of its frequency.

Conflict of interest statement.

We thereby warrant that there are not any conflicts of interests among authors and between authors and other people, institutions or organizations.

Acknowledgements.

This work was partly supported by the following grants: Ministero dell'Istruzione, dell'Università e della Ricerca 2007, 'Relationships between genetic resistance to transmissible spongiform encephalopathies and potential risk for public health and economic and biological efficiency in sheep breeds of the Piedmont Region' (P. Sacchi); Università degli Studi di Torino Ricerca scientifica finanziata 2008, 'Selezione per la resistenza genetica alla scrapie nella pecora: verifica dei risultati preliminari e previsione di possibili effetti indesiderati' (R. Rasero); and Regione Piemonte Ricerca sanitaria finalizzata 2009, 'Previsione di possibili effetti indesiderati della selezione per la resistenza genetica alla scrapie nella pecora' (R. Rasero).

References.

308 Acutis, P.L., Ru, G., Riina, M.V., Ligios, C., Maestrone, C., Fraghì, A., Carta, A.,
 309 Vaccari, G., Agrimi, U., Maroni, A., Caramelli, M., 2003. L'analisi genetica sul territorio
 310 Nazionale - risultati dell'attività del Centro di Referenza Nazionale, in: Proceedings
 311 Congress SIPAOC 'La Genetica per il controllo della Scrapie: una sfida zootecnica',
 312 Perugia (Italy), 28 November 2003.

313 Alfonso, L., Parada, A., Legarra, A., Ugarte, E., Arana, A., 2006. The effects of
 314 selective breeding against scrapie susceptibility on the genetic variability of Latxa Black-
 315 Faced sheep. *Gen. Sel. Evol.* 38, 495-511.

316 Álvarez, I., Gutiérrez, J.P., Royo, L.J., Fernández, I., Goyache, F., 2009.
 317 Quantifying diversity losses due to selection for scrapie resistance in three endangered
 318 Spanish sheep breeds using microsatellite information. *Prev. Vet. Med.* 91, 172-178.

319 Álvarez, I., Royo, L.J., Gutiérrez J.P., Fernández, I., Arranz, J.J., Goyache, F., 2007.
 320 Genetic diversity loss due to selection for scrapie resistance in the rare Spanish Xalda sheep
 321 breed. *Livest. Sci.* 111, 204-212.

322 Belt, P.B., Muilerman, I.H., Schreuder, B.E., Bos-de Ruijter, J., Gielkens, A.L.
 323 Smits, M.A., 1995. Identification of five allelic variants of the sheep PrP gene and their
 324 association with natural scrapie. *J. Gen. Virol.* 76, 509-517.

325 Bossers, A., Schreuder, B.E., Muilerman, I.H., Belt, P.B., Smits, M.A., 1996. PrP
 326 genotype contributes to determining survival times of sheep with natural scrapie. *J. Gen.*
 327 *Virol.* 77, 2660-2673.

328 Caballero, A., Toro, M.A., 2002. Analysis of genetic diversity for the management
 329 of conserved subdivided population. *Conserv. Genet.* 3, 289-299.

330 Dawson, M., Moore, R.C., Bishop, S.C., 2008. Progress and limit of PrP gene
 331 selection policy. *Vet. Res.* 39, 25-36.

332 Decreto Ministeriale 17 Dicembre 2004. Piano nazionale di selezione genetica per
 333 la resistenza alle encefalopatie spongiformi negli ovini. Gazzetta Ufficiale 51 03/03/2005.
 334 Drögemüller, C., de Vries, F., Hamann, H., Distl, O., 2004. Breeding German sheep
 335 for resistance to scrapie. Vet. Rec. 154, 257-260.
 336 European Commission, Decision 2003/100/EC of February 13, 2002, laying down
 337 minimum requirements for the establishment of breeding programmes for resistance to
 338 transmissible spongiform encephalopathies in sheep. Official Journal L041, P.0041-0045.
 339 Excoffier, L., Laval, G., Schneider, S., 2005. Arlequin ver.3.0: An integrated
 340 software package for population genetics data analysis. Evol. Bioinform. Online 1, 47-50.
 341 Falconer, D.S., Mackay, T.F.C., 1996. Introduction to Quantitative Genetics, fourth
 342 ed. Longmans Green, Harlow, Essex, UK.
 343 FAO-UNEP, 1993. World watch list for domestic animal diversity, Loftus, R.,
 344 Sherf, B. (Eds), Food and Agricultural Organization, Rome, Italy.
 345 Geldermann, H., Preuss, S., Eckert, J., Han, Y., Ollesch, K., 2003. Analysis of
 346 polymorphic microsatellites within the bovine and ovine prion protein (PRNP) genes.
 347 Anim. Genet. 34, 283-289.
 348 Goldmann, W., Baylis, M., Chihota, C., Stevenson, E., Hunter, N., 2005.
 349 Frequencies of PrP gene haplotypes in British sheep flocks and the implications for
 350 breeding programmes. J. Appl. Microbiol. 98, 1294–1302.
 351 Goudet, J., 1995. FSTAT (version 1.2): a computer program to calculate F-statistics.
 352 J. Hered. 86, 485–486.
 353 Gutiérrez, J.P., Royo, L.J., Álvarez, I. Goyache, F., 2005. MolKin v2.0: a computer
 354 program for genetic analysis of population using molecular coancestry information. J.
 355 Hered. 96,718-721.

356

357 Hunter, N., Foster, J.D., Goldmann, W., Stear, M.J., Hope, J., Bostock, C., 1996.

358 Natural scrapie in a closed flock of Cheviot sheep occurs only in specific PrP genotypes.

359 Arch. Virol. 141, 809-824.

360 Hurlbert, S.H., 1971. The non concept of species diversity: a critique and alternative

361 parameters. Ecology 52, 577-586.

362 Isler, B.J., Freking, B.A., Thallman, R.M., Heaton, M.P., Leymaster, K.A., 2006.

363 Evaluation of associations between prion haplotypes and growth, carcass, and meat quality

364 traits in a Dorset x Romanov sheep population. J. Anim. Sci. 84, 783-788.

365 Lükken, G., Brandt, H.R., Buschmann, A., Groschup, M.H., Erhardt, G., 2006.

366 Microsatellite CTSBJ12 is located distal to the ovine prion protein gene on OAR13 and is

367 not associated with scrapie susceptibility. Anim. Genet. 37, 426-427.

368 Man, W.Y.N., Nicholls, N., Woolhouse, M.E.J., Lewis, R.M., Villanueva, B., 2009.

369 Evaluating different PrP genotype selection strategies for expected severity of scrapie

370 outbreaks and genetic progress in performance in commercial sheep. Prev. Vet. Med. 91,

371 161-171.

372 Molina, A., Juárez, M., Rodero, A., 2006. Merino sheep breed's genetic resistance

373 to scrapie: genetic structure and comparison of five eradication strategies. Prev. Vet. Med.

374 75, 239-250.

375 Nei, M., 1987. Molecular evolutionary genetics, Columbia University Press, New

376 York, USA.

377 Palhière, I., Brochard, M., Verrier, É., Moazami-Goudarzi, K., Amigues, Y.,

378 Barillet, F., Bed'hom, B., Bibé, B., Bouix, J., François, D., Leymarie, C., Pantano, T.,

379 2006. Did the selection for scrapie resistance impact the genetic variability? Preliminary


380 results for French sheep breeds. CD-ROM communication 30:10. In: Proceedings of the 8th
 381 World Congress on Genetics Applied to Livestock Production, Belo Horizonte, Brasil.
 382 Palhière, I., Brochard, M., Moazami-Goudarzi, K., Laloë, D., Amigues, Y.,
 383 Bed'hom, B., Neuts, É., Leymarie, C., Pantano, T., Cribiu, E.P., Bibé, B., Verrier, É., 2008.
 384 Impact of strong selection for the PrP major gene on genetic variability of four French
 385 sheep breeds. *Gen. Sel. Evol.* 40, 663-680.
 386 Parada, A., Arana, A., Alfonso, L., 2007. Testing the equivalence of different ovine
 387 PrP genotypes for estimated breeding values. *J. Anim. Breed. Genet.* 124, 208-213.
 388 Petit, R.J., El Mousadik, A., Pons, O., 1998. Identifying populations for
 389 conservation on the basis of genetic markers. *Conserv. Biol.* 12, 844-855.
 390 Pompanon, F., Bonin, A., Bellemain, E., Taberlet, P., 2005. Genotyping errors:
 391 causes, consequences and solutions. *Nat. Rev. Genet.* 6, 847-859.
 392 Preuss, S., Kuss, A.W., He, H., Bartenschlager, H., Geldermann, H., 2005. Analysis
 393 of polymorphic PRNP microsatellite and ORF sites in German sheep breeds. *J. Anim.*
 394 *Breed. Genet.* 122 (Issue s1), 64-70.
 395 Regione Piemonte, Deliberazione della Giunta Regionale 10 ottobre 2005.
 396 Bollettino Ufficiale 42 20/10/2005.
 397 Roden, J.A., Nieuwhof, G.J., Bishop, S.C., Jones, D.A., Haresign, W., Gubbins, S.,
 398 2006. Breeding programmes for TSE resistance in British sheep. I. Assessing the impact on
 399 prion protein (PrP) genotype frequencies. *Prev. Vet. Med.* 73, 1-16.
 400 Soglia, D., Rossi, L., Cauvin, E., Citterio, C., Ferroglio, E., Maione, S., Meneguz,
 401 P.G., Spalenza, V., Rasero, R., Sacchi, P., 2010. Population genetic structure of Alpine
 402 chamois (*Rupicapra r. rupicapra*) in the Italian Alps. *Europ. J. Wild. Res.* 56, 845-854.

403 Windig, J.J., Eding, H., Moll, L., Kaal, L., 2004. Effects on inbreeding of different
404 strategies aimed at eliminating scrapie sensitivity alleles in rare sheep breeds in The
405 Netherland. Anim. Sci. 79, 11-20.

406 Wiśniewska, E., Mroczkowski, S., 2010. Prion protein (PrP) gene polymorphism
407 and simulation study of breeding oriented to scrapie resistance in Polish Merino and Polish
408 Mountain Sheep. Sci. Ann. Pol. Soc. Anim. Prod. 6, 115-123.

409

410 Figure 1 Differences in allele arrangements (F_{ST}) between the before-2005 and after-2005 cohorts of
411 rams as a function of distance of any OAR13 microsatellite from PRNP (deviations are in Mb). Grey
412 circles show the OAR13 microsatellites outside PRNP, empty circles show the OAR13 microsatellites
413 within PRNP, the black circle points at PRNP. All F_{ST} values above the dotted line are statistically
414 significant.



1 **Table 1**

2 Location and values of F_{IS} for the OAR13 and neutral microsatellites.

3

OAR13 microsatellites				Neutral microsatellites			
	Dist. ^a	Before-2005	After-2005		OAR ^b	Before-2005	After-2005
BMC1222	-35.1	+0.080 n.s.	+0.089 n.s.	CSRD247	14	+0.031 n.s.	+0.197 **
MCM152	-18.5	-0.081 n.s.	-0.028 n.s.	D5S2	5	+0.058 n.s.	+0.243 **
ILSTS59	-17.0	-0.105 n.s.	+0.025 n.s.	HSC	9	+0.068 n.s.	+0.142 **
HUJ616	-8.2	+0.065 n.s.	-0.115 *	INRA23	1	+0.084 n.s.	+0.102 n.s.
URB58	-1.4	-0.089 n.s.	-0.009 n.s.	INRA5	10	+0.429 ***	+0.212 ***
BMS1669	-0.6	-0.059 n.s.	+0.096 n.s.	INRA63	14	+0.072 n.s.	-0.006 n.s.
PRNPS04	0	-0.085 n.s.	-0.077 n.s.	MAF65	15	+0.010 n.s.	+0.011 n.s.
PRNPS05	0	-0.248 ***	-0.286 ***	MCM527	5	+0.047 n.s.	+0.064 n.s.
PRNPS11	0	-0.045 n.s.	+0.290 *	OarCP49	17	-0.070 n.s.	+0.017 n.s.
PRNPS15	0	+0.166 n.s.	+0.018 n.s.	OarFCB11	2	+0.005 n.s.	+0.130 *
PRNPS24	0	+0.166 n.s.	-0.128 n.s.	OarFCB20	2	+0.062 n.s.	+0.121 n.s.
CTSB12	+21.0	-0.025 n.s.	+0.002 n.s.	OarFCB304	19	-0.088 n.s.	-0.051 n.s.
MMP9	+27.9	+0.074 n.s.	+0.259 ***				
BMS2319	+32.5	+0.110 *	+0.399 ***				

4

5 ^a Distance of any locus from PRNP (deviation in Mb, Ovine version 2.0 Genome Assembly map

6 provided by the International Sheep Genomic Consortium, [http://www.livestockgenomics.csiro.au/cgi-](http://www.livestockgenomics.csiro.au/cgi-bin/gbrowse/oarv2.0/)
7 [bin/gbrowse/oarv2.0/](http://www.livestockgenomics.csiro.au/cgi-bin/gbrowse/oarv2.0/)).

8 ^b Chromosome location.

9 n.s. not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

10

11 **Table 2**

12 Summary of genetic variability in the before-2005 and after-2005 cohorts of rams at PRNP locus.

13

	Before-2005	After-2005
Allele frequencies		
ARR	0.289	0.577
ARQ	0.570	0.354
AHQ	0.028	0.038
ARH	0.014	0.000
VRQ	0.099	0.031
Genotype frequencies		
ARR/ ARR	0.028	0.354
ARR/ AHQ	0.014	0.046
AHQ/ AHQ	0	0
ARR/ ARQ	0.437	0.385
ARR/ ARH	0.014	0
ARQ/ AHQ	0.042	0.031
AHQ/ ARH	0	0
ARR/ VRQ	0.056	0.015
ARQ/ ARQ	0.282	0.123
ARQ/ ARH	0	0
AHQ/ VRQ	0	0
ARH/ ARH	0	0
ARQ/ VRQ	0.099	0.046
ARH/ VRQ	0.014	0
VRQ/ VRQ	0.014	0
Summary of genetic variability		
GD	0.581	0.540
F _{IS}	-0.158 *	+0.038 n.s.
A	5.0	4.0

14

15 n.s. not significant; * P < 0.05.

16

17 **Table 3**

18 F_{ST} values between the before-2005 and after-2005 cohorts of rams for PRNP and on the whole for
19 OAR13 microsatellite markers and neutral markers.

PRNP	0.101 ***
OAR13 markers within PRNP	0.021 (0.011) **
OAR13 markers outside PRNP	0.011 (0.003) ***
Neutral markers	0.005 (0.002) n.s.

20
21 Standard error in parenthesis.

22 n.s. not significant, ** $P < 0.01$; *** $P < 0.001$.

23

24

Table 4

GD, F_{IS} , PIC, A, and A_g values in the before-2005 and after-2005 cohorts of rams at three clusters of microsatellite markers on different locations.

	GD	F_{IS} ^a	PIC	A	A_g ^b
OAR13 markers within PRNP					
Before-2005	0.410	−0.018 n.s.	0.293	3.0	3.0
After-2005	0.302	−0.087 n.s.	0.215	3.0	3.0
OAR13 markers outside PRNP					
Before-2005	0.761	0	0.589	8.3	8.1
After-2005	0.755	+0.075 *	0.589	8.2	8.1
Neutral markers					
Before-2005	0.739	+0.065 *	0.605	8.8	8.6
After-2005	0.730	+0.104 *	0.592	9.3	9.2

^a Based on 10000, 18000, and 24000 randomizations for the three different locations, respectively. n.s. not significant; * significant at 0.05 adjusted nominal level.

^b Adjusted for $g = 128, 118$, and 124 for the three different locations, respectively.